

Recognizing Idiopathic Condylar Resorption: A Case-Based Review

Ismail Yardimcioglu, MD¹; Jordan Hughes, MD¹; Neil Tishkoff, MD²

1) Yale New Haven Health – Bridgeport Hospital, Department of Radiology

2) Yale University School of Medicine, Radiology & Biomedical Imaging

Outline

- Disclosures
- Teaching Points
- Introduction
- Cases
- Discussion

Disclosures

- None

Teaching Points

- Describe the epidemiology and clinical features of idiopathic condylar resorption (ICR)
- Recognize the importance of ICR
- Review the imaging findings
- Differentiate ICR from other causes of condylar resorption
- Highlight the role of radiologists in early detection and prevention of future complications

Introduction

- Idiopathic condylar resorption, also known as progressive condylar resorption is an uncommon yet clinically significant condition characterized by non-inflammatory progressive loss of the mandibular condyle.
- The condition results from progressive lysis and repair of the articular cartilage and underlying condylar bone. This remodeling process leads to condylar height loss which can cause mandibular retrognathia and malocclusion.

Introduction

- It occurs predominantly in adolescent females ($\approx 90\%$), with hormonal influences thought to play a role in its pathogenesis and rarely seen after age of 23.
- Presenting symptoms may include jaw pain, joint clicking, limited mandibular range of motion, malocclusion, facial asymmetry, and altered bite efficiency.

Introduction

- Primary: Idiopathic condylar resorption which occurs without identifiable systemic or local pathology and is typically bilateral.
- Secondary: Causes include trauma, prior temporomandibular joint (TMJ) surgery, corticosteroid use, or systemic autoimmune diseases.

Introduction

- Primary idiopathic condylar resorption is a diagnosis of exclusion, with juvenile idiopathic arthritis (JIA) the most significant differential diagnosis.
- Accurate diagnosis relies on clinical correlation and imaging findings.
- Radiologists are critical in identifying early imaging findings of the disease process, establishing diagnosis, evaluating disease extent and progression, assessing stability and guiding management decisions.

Introduction

- Both CT and MRI are essential—CT for detailed assessment of osseous changes and MRI for evaluating soft tissue, disc morphology, bone marrow signal and disease activity.
- Typical imaging findings include:
 - CT: Erosion and flattening of the superior condylar surface, loss of condylar volume, shallow glenoid fossa, and flattened articular eminence.
 - MRI: Characteristic anterior displacement of the disc without recapture, condylar flattening/erosion, and reduced condylar volume. Some authors also report bone marrow edema and peri-condylar contrast enhancement during the active phase.

Introduction

- Management strategies vary depending on disease activity and progression.
 - Non-surgical approaches may include NSAIDs, splint therapy and observation.
 - Surgical options include disc repositioning, joint reconstruction, orthognathic procedures or temporomandibular joint replacement. Surgical procedures are typically deferred until the resorptive process has stabilized.
- Early diagnosis and regular follow-up are as crucial as definitive treatment.

Cases

Case 1:

A 12-year-old female with a history of retrognathia developed new occlusal changes after completing Stage I orthodontic treatment for Class III malocclusion.

She has no prior medical history, trauma, or surgical interventions. The patient denies temporomandibular joint (TMJ) pain, clicking, or other joint-related symptoms.

On examination, a new open bite discrepancy was identified without TMJ tenderness or audible sounds.

Laboratory evaluation was unremarkable.





Fig.1: Flattening of the right mandibular condyle (A: Closed mouth coronal proton density), and abnormal enhancement surrounding the temporomandibular joint and mandibular condyle (B: Closed mouth coronal T1 fat-sat postcontrast).




Fig.2: Sagittal proton density images (A: Closed mouth and B: Open mouth) demonstrate thinning of the articular disc as well as flattening/resorption of the left mandibular condyle, decreased range of motion. C: Coronal post contrast T1 fat-sat image reveals abnormal enhancement surrounding the left temporomandibular joint.




Fig.3: A: Sagittal and B: Coronal CT images demonstrate flattening and irregularity/small erosions, of the right mandibular condyle and shallow glenoid fossa.




Fig.4: A: Sagittal and B: Coronal CT images demonstrate flattening, microcystic erosions and irregularity of the left mandibular condyle.

- Imaging findings are most consistent with idiopathic condylar resorption (ICR); however, other inflammatory conditions remain in the differential given the presence of contrast enhancement.
- Although the rheumatologic workup was unremarkable, excluding systemic inflammatory disorders such as juvenile idiopathic arthritis (JIA), the patient will be started on an anti-TNF- α agent as empiric therapy.
- Plastic surgery and oral maxillofacial surgery teams have deferred intervention at this time but anticipate possible surgical management if her symptoms or imaging findings worsen.

Case 2:

A 17-year-old female with no past medical history presents for a three-year history of gradual-onset jaw pain and intermittent clicking.

Rheumatologic evaluation was negative, effectively excluding secondary causes.

Initial MRI demonstrated bilateral condylar flattening and irregularity within shallow glenoid fossae, accompanied by restricted mandibular motion.




Fig.5: A: Closed mouth sagittal T2 and B: Closed mouth sagittal proton-density images demonstrate abnormal appearance of the right mandibular condyle which is seated within abnormally shallow glenoid fossa.




Fig.6: 2 years later. Right sagittal closed mouth views (A: T2 fat-sat and B: T1) demonstrate erosive changes of the mandibular condyle in addition to flattening of the condyle and glenoid fossa. The articular disc is also markedly degenerated and only faintly identified. Overall findings worsened in the interim.




Fig.7: Left sagittal closed mouth views A: T2 fat-sat and B: T1 demonstrate flattening of the mandibular condyle within an abnormally shallow glenoid fossa. The articular disc is also markedly degenerated and only faintly identified.

She has been monitored with annual MRI for 3 years, which has shown progressive condylar resorption.

Surgical intervention is now being considered as the patient approaches skeletal maturity.

Conclusion

- Idiopathic condylar resorption is a rare but important cause of temporomandibular joint dysfunction, predominantly affecting adolescent females and typically bilateral.
- Early recognition of ICR is essential to prevent long-term complications.

Conclusion

- Characteristic imaging findings include but not limited to condylar flattening, microcystic bony changes, shallow glenoid fossa, disc displacement and atrophy, and bone marrow edema.
- While ICR is generally non-inflammatory and contrast enhancement is uncommon, it may be observed in cases with active resorption.

Conclusion

- Differentiating idiopathic condylar resorption from juvenile idiopathic arthritis is based on both clinical and imaging findings.
 - Clinically, JIA typically presents with multiple joint involvement and positive inflammatory markers on laboratory evaluation, whereas ICR is usually isolated to the temporomandibular joints with normal laboratory results.
 - On imaging, JIA often demonstrates synovial proliferation, soft tissue edema, and enhancement, along with punched-out erosions and evidence of active inflammation. In contrast to ICR, disc displacement is not commonly observed in JIA.

Conclusion

- Radiologists should maintain a high index of suspicion in the appropriate clinical setting and play a central role in early detection, guiding multidisciplinary approach and preventing long-term complications.
- Clinical clues such as jaw pain, restricted joint motion, or new occlusal changes should raise suspicion and alert the interpreting radiologist to carefully evaluate the temporomandibular joints.

Conclusion

- Management of ICR depends on patient's age, the extent of temporomandibular joint (TMJ) involvement and the severity of symptoms and includes:
 - Conservative therapies such as analgesics, physical therapy and occlusal splints
 - Surgical management such as disc repositioning (discopexy), corrective orthognathic surgery, surgical repositioning of TMJ disc and stabilization or, in advanced cases, total TMJ replacement with prosthesis.

References

1. Larheim TA, Hol C, Løseth G, Arvidsson LZ. Temporomandibular joint pathologies: pictorial review. *Br J Radiol*. 2024 Jan 23;97(1153):53-67. doi: 10.1093/bjr/tqad021. PMID: 38263820; PMCID: PMC11027251.
2. Zarour CC, Robinson C, Mian A, Al-Hameed M, Vempala M. Idiopathic Mandibular Condyle Resorption. *Cureus*. 2020 Nov 6;12(11):e11365. doi: 10.7759/cureus.11365. PMID: 33304698; PMCID: PMC7721076.
3. Prado BN, Pereira LC, Prado JN. Aggressive idiopathic condylar reabsorption after orthognathic surgery: A complex Imaging diagnosis. *Radiol Case Rep*. 2025 Mar 23;20(6):2940-2945. doi: 10.1016/j.radcr.2025.02.098. PMID: 40207031; PMCID: PMC11979931.
4. Wolford LM. Idiopathic condylar resorption of the temporomandibular joint in teenage girls (cheerleaders syndrome). *Proc (Bayl Univ Med Cent)*. 2001 Jul;14(3):246-52. doi: 10.1080/08998280.2001.11927772. PMID: 16369629; PMCID: PMC1305829.
5. Tanaka E, Mercuri LG. Current Status of the Management of Idiopathic Condylar Resorption/Progressive Condylar Resorption-A Scoping Review. *J Clin Med*. 2024 Jul 5;13(13):3951. doi: 10.3390/jcm13133951. PMID: 38999512; PMCID: PMC11242904.
6. McLoughlin PM, Hopper C, Bowley NB. Hyperplasia of the mandibular coronoid process: an analysis of 31 cases and a review of the literature. *J Oral Maxillofac Surg*. 1995 Mar;53(3):250-5. doi: 10.1016/0278-2391(95)90219-8. PMID: 7861274.

Questions

- Please e-mail with any questions
ismailyardimcioglu@gmail.com